

Statistical bulletin

# Coronavirus (COVID-19) Infection Survey, antibody and vaccination data, UK: 8 December 2021

Antibody and vaccination data by UK country and regions in England from the Coronavirus (COVID-19) Infection Survey. This analysis has been produced in partnership with University of Oxford, University of Manchester, UK Health Security Agency and Wellcome Trust. This study is jointly led by the ONS and the Department for Health and Social Care (DHSC) working with the University of Oxford and Lighthouse Laboratories to collect and test samples.

Contact:  
Rhiannon Yapp and Byron  
Davies  
infection.survey.analysis@ons.  
gov.uk  
+44 1633 560499

Release date:  
8 December 2021

Next release:  
23 December 2021

## Table of contents

1. [Main points](#)
2. [Understanding antibodies and immunity](#)
3. [Percentages of adults testing positive for COVID-19 antibodies and vaccinated against COVID-19 in England, Wales, Northern Ireland and Scotland](#)
4. [Percentages of adults testing positive for COVID-19 antibodies and vaccinated against COVID-19 by regions in England](#)
5. [Percentages of adults testing positive for COVID-19 antibodies and vaccinated against COVID-19 by grouped age in England, Wales, Northern Ireland and Scotland](#)
6. [Percentage of adults testing positive for COVID-19 antibodies by single year of age in England, Wales, Northern Ireland and Scotland](#)
7. [Coronavirus \(COVID-19\) Infection Survey data](#)
8. [Collaboration](#)
9. [Glossary](#)
10. [Measuring the data](#)
11. [Strengths and limitations](#)
12. [Related links](#)

# 1 . Main points

The following statistics are the percentage of the adult population aged 16 and over, that would have tested positive on a blood test for antibodies against SARS-CoV-2, the specific virus that causes coronavirus (COVID-19). This suggests that they have had the infection in the past or have been vaccinated.

- In England, it is estimated that 95.3% of the adult population (95% credible interval: 94.5% to 96.0%) would have tested positive for antibodies against SARS-CoV-2 in the week beginning 15 November 2021.
- In Wales, it is estimated that 93.9% of the adult population (95% credible interval: 92.7% to 95.0%) would have tested positive for antibodies against SARS-CoV-2 in the week beginning 15 November 2021.
- In Northern Ireland, it is estimated that 91.6% of the adult population (95% credible interval: 87.9% to 93.5%) would have tested positive for antibodies against SARS-CoV-2 in the week beginning 15 November 2021.
- In Scotland, it is estimated that 95.0% of the adult population (95% credible interval: 94.0% to 95.9%) would have tested positive for antibodies against SARS-CoV-2 in the week beginning 15 November 2021.
- The percentage of the adult population who would have tested positive for antibodies against SARS-CoV-2 has increased in those aged 65 years and over across the UK, likely as a result of the vaccination booster programme.

## About this bulletin

This bulletin presents the latest estimates of the number of adults who would test positive for antibodies to SARS-CoV-2 for England, Wales, Northern Ireland and Scotland based on findings from the Coronavirus (COVID-19) Infection Survey in the UK. People who test positive for antibodies are those who have experienced previous infection and/or vaccination. We include estimates from our survey on the percentage of those aged 16 and over who reported they have received one or more doses of a vaccine against SARS-CoV-2 (including booster doses).

## Data in this bulletin

The analysis on antibodies in this bulletin is based on blood test results taken from a randomly selected subsample of individuals aged 16 years and over who live in private households. The survey excludes those in hospitals, care homes and/or other communal establishments. The blood samples are used to test for antibodies against SARS-CoV-2.

We also present data on the percentage of people aged 16 years and over who report they have received one or more doses of a COVID-19 vaccine since 14 December 2020, and two or more doses of a COVID-19 vaccine since 15 February 2021.

Our antibodies and vaccination estimates are based on the data we collect from people visited in the Coronavirus (COVID-19) Infection Survey (CIS). We present weekly modelled antibody estimates and vaccine estimates for adults by country, and grouped age for England, Wales, Northern Ireland and Scotland, as well as antibody estimates by regions in England. Further information on our method to model antibodies and vaccinations can be found in [our methods article](#).

There is a clear pattern between vaccination and testing positive for COVID-19 antibodies but the detection of antibodies alone is not a precise measure of the [immunity protection given by vaccination](#). Further information is available on [antibody levels post vaccination](#) from research partners at the University of Oxford.

Modelled vaccine estimates are produced to provide context alongside our antibodies estimates and do not replace the [official government figures on vaccinations](#), which are a more precise count of total vaccines issued. While we would expect the overall trend of our estimated number of people who have received vaccines to increase, it is possible that in some weeks, the estimate may remain the same or decrease as a result of sampling variability (for example, we may have a lower number of participants recording a vaccination in the latest week compared with an earlier week).

National Immunisation Management System (NIMS) administrative data are used to validate CIS self-reported records of vaccination for England. The equivalent of this is currently not included for other countries, meaning the estimates for Wales, Scotland and Northern Ireland are produced just from CIS self-reported records of vaccination.

Our vaccination estimates are explained in more detail in [Section 10: Measuring the data](#).

## 2 . Understanding antibodies and immunity

Antibody positivity is defined by having a fixed concentration of antibodies in the blood. A negative test result occurs if there are no antibodies, or if antibody levels are too low to reach a threshold at the time of testing. The threshold we use is 42 nanograms per millilitre (ng/ml). This is the threshold that the test is CE marked against providing greater than 99% [sensitivity and specificity](#). Following infection or vaccination, antibody levels can vary and sometimes increase but may still be below the level identified as positive in our test and other tests, and so give a negative result.

Most people who are vaccinated will retain a higher antibody level than before vaccination even after dropping below the standard threshold value. A negative result does not mean that the antibody level is at zero, nor that a person has no protection against COVID-19, as an immune response does not rely on the presence of antibodies alone.

There are other parts of the immune system that will offer protection, for example, a person's T-cell response. This will not be detected by blood tests for antibodies. [A person's immune response is affected by a number of factors](#), including health conditions and age.

Equally, antibody levels are expected to decrease over time irrespective of vaccination or natural infection, especially when exposure to the virus is reduced. This is because our bodies stop making antibodies when they are not needed.

Our [blog on antibodies and immunity](#) gives further information on the link between antibodies and immunity and the vaccine programme. [Our blog on vaccine effectiveness](#) provides information on the effectiveness of vaccinations against Alpha and Delta variants, which is based upon the research conducted by partners from the University of Oxford.

Our [methodology article](#) provides further information around the survey design, how we process data, and how data are analysed. The [study protocol](#) specifies the research for the study. The [Coronavirus \(COVID-19\) Infection Survey QMI](#) details the strength and limitations of the data.

### 3 . Percentages of adults testing positive for COVID-19 antibodies and vaccinated against COVID-19 in England, Wales, Northern Ireland and Scotland

In the week beginning 15 November 2021, the percentage of adults testing positive for coronavirus (COVID-19) antibodies remained high across the UK.

Table 1: Estimated percentage of adults who would have tested positive for SARS-CoV-2 antibodies from a blood sample, week beginning 15 November 2021, UK countries

Country	Estimated % of adults who would have tested positive 95% credible interval for COVID-19 antibodies		
		Lower	Upper
<b>England</b>	95.3	94.5	96.0
<b>Wales</b>	93.9	92.7	95.0
<b>Northern Ireland</b>	91.6	87.9	93.5
<b>Scotland</b>	95.0	94.0	95.9

Source: Office for National Statistics – Coronavirus (COVID-19) Infection Survey

#### Notes

1. All estimates are subject to uncertainty given that a sample is only part of the wider population. The model used to provide these estimates is a Bayesian model: these provide 95% credible intervals.

In the week beginning 15 November 2021, our estimates suggest the percentage of adults in England who reported they had received at least one dose of a COVID-19 vaccine remained high, at 96.5% (95% credible interval: 95.1% to 97.4%). The estimated percentage of adults in England who had received two or more doses of a COVID-19 vaccine also remained high, with 91.8% (95% credible interval: 88.3% to 94.3%) of adults receiving two or more doses in the week beginning 15 November 2021. These vaccination estimates will differ from official figures as explained in [Section 1](#).

Modelled vaccination estimates for Wales, Northern Ireland, and Scotland require additional quality assurance and are not included in our bulletin this time. We will reintroduce our vaccine estimates as soon as possible for these countries.

#### Figure 1: The percentage of adults testing positive for COVID-19 antibodies remained high across UK countries in the week beginning 15 November 2021

Modelled percentage of adults: who tested positive for antibodies to SARS-CoV-2, UK Countries, and who have received one or more doses of a COVID-19 vaccine; and two or more doses of a COVID-19 vaccine, England, 7 December 2020 to 20 November 2021

#### Notes:

1. All results are provisional and subject to revision.
2. These statistics refer to antibody tests and reported vaccinations for individuals living in private households. These figures exclude individuals in hospitals, care homes and/or other communal establishments.
3. All estimates are subject to uncertainty, given that a sample is only part of the wider population. A credible interval gives an indication of the uncertainty of an estimate from data analysis.
4. The denominators used for both antibodies and vaccinations are the total people in the sample at that particular time point, then post-stratified by the mid-year population estimate.
5. Our estimates of vaccination are provided for context alongside our antibodies estimates but are likely to be different from the official figures. The [daily official government figures](#) provide the recorded actual numbers of vaccines against SARS-CoV-2 issued.
6. Modelled vaccination estimates for Wales, Northern Ireland and Scotland have not been included this time on account of additional quality assurance required

[.xlsx](#)

## **4 . Percentages of adults testing positive for COVID-19 antibodies and vaccinated against COVID-19 by regions in England**

The percentage of adults testing positive for coronavirus (COVID-19) antibodies and the percentage of adults who have received at least one vaccine remained at high levels across all regions of England in the week beginning 15 November 2021. Regional data can be found in our [Coronavirus \(COVID-19\) Infection Survey: antibody and vaccination dataset](#).

## 5 . Percentages of adults testing positive for COVID-19 antibodies and vaccinated against COVID-19 by grouped age in England, Wales, Northern Ireland and Scotland

In the week beginning 15 November 2021, the percentage of adults testing positive for coronavirus (COVID-19) antibodies remained high across all age groups in England, Wales, Northern Ireland and Scotland. There were increases in all age groups over recent weeks in England, Wales and Scotland and in most age groups in Northern Ireland. There has been an increased antibody positivity in those aged 65 years and over across the UK since early October 2021, likely as a result of the vaccination booster programme.

It is important to note that antibody positivity is defined by a fixed concentration of antibodies in the blood. Most people who are vaccinated will retain higher antibody levels than before vaccination but may have antibodies below the threshold used at the time of testing. This does not mean that these people have no protection against new infection. More information on antibodies and immunity can be found in [our blog](#) and in [Section 2](#) of this bulletin.

The percentage of adults in England who reported they have received at least one dose of a COVID-19 vaccine is lowest in the younger age groups but still remains high. In England, the estimated percentage of those aged 16 to 24 years who have received one or more doses of a COVID-19 vaccine was 86.8% (95% credible interval: 79.3% to 91.5%); this has increased sharply since the end of June 2021. These estimates of vaccination are based on reported vaccination status in our survey and are likely to be different from the [official figures](#) for the UK.

In the data used to produce estimates for Wales, Northern Ireland and Scotland, the number of people sampled who tested positive for antibodies to SARS-CoV-2 or who have been vaccinated is low compared with England. This means there is a higher degree of uncertainty in estimates for these nations when our analysis splits the sample into smaller groups (for example, age groups) as indicated by larger credible intervals.

Modelled antibody estimates and modelled vaccine estimates relate to the adult population (aged 16 years and over) who live in private households. The true figure among the older age groups in the population may be different. In England, an estimated [90% of people aged 80 years and over live in private households](#) and 10% live in other communal establishments such as care homes.

### **Figure 2: The percentage of adults testing positive for COVID-19 antibodies remained high and increased for all age groups across the UK in the week beginning 15 November 2021**

Modelled percentage of adults: who tested positive for antibodies to SARS-CoV-2, UK, who have received one or more doses of a COVID-19 vaccine, and those who have received two or more doses of a COVID-19 vaccine, by grouped age, England, 7 December 2020 to 20 November 2021

#### **Notes:**

1. All results are provisional and subject to revision.
2. These statistics refer to antibody tests and reported vaccinations for individuals living in private households. These figures exclude individuals in hospitals, care homes and/or other communal establishments.
3. In Northern Ireland, the number of people sampled is low compared with England, Wales and Scotland; therefore, people aged 50 to 69 years are included in the same age group, and those aged 70 years and over are included in the same age group.
4. All estimates are subject to uncertainty, given that a sample is only part of the wider population. A credible interval gives an indication of the uncertainty of an estimate from data analysis.
5. The denominators used for both antibodies and vaccinations are the total people in the sample at that particular time point, then post-stratified by the mid-year population estimate
6. Our estimates of vaccination are provided for context alongside our antibodies estimates but are likely to be different from the official figures. The [daily official government figures](#) provide the recorded actual numbers of vaccines against SARS-CoV-2 issued.
7. Modelled vaccination estimates for Wales, Northern Ireland and Scotland have not been included this time on account of additional quality assurance required.

[.xlsx](#)

## 6 . Percentage of adults testing positive for COVID-19 antibodies by single year of age in England, Wales, Northern Ireland and Scotland

The analysis in this section presents modelled daily estimates of coronavirus (COVID-19) antibody positivity by single year of age for England, Wales, Northern Ireland and Scotland separately, between 10 October and 20 November 2021. The modelled data in this section are produced using a different method to the weekly modelled estimates presented in [Section 5](#) and so cannot be compared. Each data point represents a modelled estimate of antibody positivity for a particular day.

This more granular analysis shows a similar pattern to our grouped age analysis, with antibody positivity remaining high across the UK. In recent weeks, antibody positivity rates have increased for those aged 70 years and over in England and those aged 65 years and over in Wales, Scotland and Northern Ireland.

### Figure 3: The percentages testing positive for COVID-19 antibodies by single year of age in England, Wales, Northern Ireland and Scotland

Modelled percentage of adults testing positive for antibodies to SARS-CoV-2, by single year of age, UK countries, 10 October to 20 November 2021

#### Notes:

1. All results are provisional and subject to revision.
2. These statistics refer to antibody tests reported in the community, by which we mean private households. These figures exclude individuals in hospitals, care homes and/or other communal establishments.
3. The method used to generate the data differs from the modelled weekly estimates of antibody positivity by age and so is not comparable.

[.xlsx](#)

## 7 . Coronavirus (COVID-19) Infection Survey data

[Coronavirus \(COVID-19\) antibody and vaccination data, UK](#)

Dataset | Released 8 December 2021

Antibody and vaccination data by UK country and regions in England from the Coronavirus (COVID-19) Infection Survey. This analysis has been produced in partnership with the University of Oxford, University of Manchester, UK Health Security Agency and Wellcome Trust. This study is jointly led by the ONS and the Department for Health and Social Care (DHSC) working with the University of Oxford and Lighthouse Laboratory to collect and test samples.

## 8 . Collaboration



UK Health  
Security  
Agency



The Coronavirus (COVID-19) Infection Survey analysis was produced by the Office for National Statistics (ONS) in collaboration with our research partners at the University of Oxford, the University of Manchester, UK Health Security Agency and Wellcome Trust. Of particular note are:

- Sarah Walker - University of Oxford, Nuffield Department for Medicine: Professor of Medical Statistics and Epidemiology and Study Chief Investigator
- Koen Pouwels - University of Oxford, Health Economics Research Centre, Nuffield Department of Population Health: Senior Researcher in Biostatistics and Health Economics
- Thomas House - University of Manchester, Department of Mathematics: Reader in Mathematical Statistics

## 9 . Glossary

## Antibodies

We measure the presence of antibodies in people who live in private households to understand who has had coronavirus (COVID-19) in the past, and the impact of vaccinations. It takes between two and three weeks after infection or vaccination for the body to make enough antibodies to fight the infection. Antibodies can help prevent individuals from getting the same infection again. If they do get infected, people with antibodies are less likely to have severe symptoms. Once infected or vaccinated, antibodies remain in the blood at low levels and can decline over time. The length of time antibodies remain at detectable levels in the blood is not fully known.

## SARS-CoV-2

This is the scientific name given to the specific virus that causes COVID-19.

## Confidence interval

A confidence interval gives an indication of the degree of uncertainty of an estimate, showing the precision of a sample estimate. The 95% confidence intervals are calculated so that if we repeated the study many times, 95% of the time the true unknown value would lie between the lower and upper confidence limits. A wider interval indicates more uncertainty in the estimate. Overlapping confidence intervals indicate that there may not be a true difference between two estimates.

For more information, see our methodology page on [statistical uncertainty](#).

## Credible interval

A credible interval gives an indication of the uncertainty of an estimate from data analysis. A 95% credible interval is calculated so that there is a 95% probability of the true value lying in the interval.

# 10 . Measuring the data

## Reference dates

We produce weekly modelled estimates using standard calendar weeks starting Monday. To provide the most timely and accurate estimates possible for antibody positivity, the model will include data for the first four to seven days of the most recent week available, depending on the availability of test results. The antibody estimates for the most recent week in this publication includes data from 15 to 20 November 2021. The vaccination estimates for the most recent week for England in this publication includes data from 8 to 14 November 2021.

More information on [measuring the data](#) is available in the Coronavirus (COVID-19) Infection Survey statistical bulletin.

Our [methodology article](#) provides further information around the survey design, how we process data, and how data are analysed. The [Quality and Methodology Information](#) explains the strengths and limitations of the data, methods used, and data uses and users.

## Vaccination estimates

While the [daily official government figures](#) provide the recorded actual numbers of vaccines against SARS-CoV-2 issued, our vaccination estimates are likely to be different from the official figures. This is because they are estimates based on a sample survey of reported vaccine status and are provided for context alongside our antibodies estimates. We control for the effect of ethnicity by post-stratifying our analysis by White and other ethnic groups, rather than individual ethnicities, because of our current sample size. This could result in differences between our survey estimates and the government figures in the numbers of vaccines received for some ethnic minority groups.

Importantly, our survey collects information from the population living in private households and does not include people living in communal establishments such as care homes, hospitals or prisons.

The value of showing our estimates of vaccines alongside our estimates of people testing positive for antibodies is to illustrate the relationship between the two.

Differences between official figures and the estimates from this survey differ in scale across each of the four UK nations (some survey estimates are closer to the official reported figures than others) because of differences in reporting dates and the inclusion of National Immunisation Management System (NIMS) data for England. In addition, our sampling method for Northern Ireland is different to the other nations, inviting only people who have previously participated in a Northern Ireland Statistics and Research Agency (NISRA) survey, which could result in a sample of individuals who are more likely to get vaccinated. This should be taken into consideration if comparing vaccine and antibody estimates across the four nations, as vaccine status and antibody positivity are related.

In addition, as our analysis develops, our survey-based estimates will enable possible future analysis of people who have received a vaccine with other characteristics collected in the survey.

## 11 . Strengths and limitations

More information on strengths and limitations of the data is available in [Coronavirus \(COVID-19\) Infection Survey QMI](#) and in the [Coronavirus \(COVID-19\) Infection Survey statistical bulletin](#).

## 12 . Related links

### [Coronavirus \(COVID-19\) Infection Survey, UK](#)

Bulletin | Updated weekly

Estimates for England, Wales, Northern Ireland and Scotland. This survey is being delivered in partnership with the University of Oxford, University of Manchester, UK Health Security Agency and Wellcome Trust.

### [Coronavirus \(COVID-19\) Infection Survey: characteristics of people testing positive for COVID-19 in England](#)

Bulletin | Updated fortnightly

Characteristics of people testing positive for COVID-19 from the Coronavirus (COVID-19) Infection Survey, including antibody data by UK country, and region and occupation for England. Antibodies data published before 3 February 2021 are available in this series.

### [Coronavirus \(COVID-19\) Infection Survey technical article: Analysis of characteristics associated with vaccination uptake](#)

Technical article | Released 15 November 2021

Analysis of populations in the UK by likelihood of being vaccinated against COVID-19 using the COVID-19 Infection Survey. This survey is being delivered in partnership with University of Oxford, University of Manchester, UK Health Security Agency and Wellcome Trust.

### [Coronavirus \(COVID-19\) Infection Survey Technical Article: Impact of vaccination on testing positive in the UK: October 2021](#)

Technical article | Released 18 October 2021

The reduction in risk of testing positive for COVID-19 associated with vaccination overall and by different vaccine types using data from the Coronavirus (COVID-19) Infection Survey. Two time periods were analysed; when the Alpha variant was dominant in the UK (1 December 2020 to 16 May 2021), and when the Delta variant was dominant (17 May to 14 August 2021).

### [Coronavirus \(COVID-19\) Infection Survey QMI](#)

QMI | Released 16 July 2021

Quality and Methodology Information for the Coronavirus (COVID-19) Infection Survey (CIS), detailing the strengths and limitations of the data, methods used, and data and users.