

Statistical bulletin

# Coronavirus (COVID-19) Infection Survey, UK: 14 April 2022

Estimates for England, Wales, Northern Ireland and Scotland. This survey is being delivered in partnership with University of Oxford, University of Manchester, UK Health Security Agency and Wellcome Trust. This study is jointly led by the Office for National Statistics (ONS) and the Department for Health and Social Care (DHSC) working with the University of Oxford and Lighthouse laboratory to collect and test samples.

Contact:  
Rhiannon Yapp and Mike  
Bracher  
infection.survey.analysis@ons.  
gov.uk  
+44 1633 560499

Release date:  
14 April 2022

Next release:  
22 April 2022

## Table of contents

1. [Main points](#)
2. [Percentage of people who had COVID-19 in UK countries](#)
3. [Sub-national analysis of the percentage of people who had COVID-19](#)
4. [Age analysis of the percentage of people who had COVID-19](#)
5. [Number of new COVID-19 infections in UK countries](#)
6. [Analysis of viral load and variants of COVID-19](#)
7. [Test sensitivity and specificity](#)
8. [Coronavirus \(COVID-19\) Infection Survey data](#)
9. [Collaboration](#)
10. [Glossary](#)
11. [Measuring the data](#)
12. [Strengths and limitations](#)
13. [Related links](#)

# 1 . Main points

- In England, the percentage of people testing positive for coronavirus (COVID-19) decreased in the week ending 9 April 2022; we estimate that 3,773,800 people in England had COVID-19 (95% credible interval: 3,662,100 to 3,888,000), equating to 6.92% of the population or around 1 in 14 people.
- In Wales, the percentage of people testing positive for COVID-19 remained high in the most recent week (up to 9 April 2022); we estimate that 231,900 people in Wales had COVID-19 (95% credible interval: 206,200 to 259,800), equating to 7.63% of the population or around 1 in 13 people.
- In Northern Ireland, the trend in the percentage of people testing positive for COVID-19 decreased in the week ending 9 April 2022; we estimate that 95,900 people in Northern Ireland had COVID-19 (95% credible interval: 78,800 to 114,000), equating to 5.23% of the population or around 1 in 19 people.
- In Scotland, the percentage of people testing positive for COVID-19 continued to decrease in the week ending 9 April 2022; we estimate that 314,800 people in Scotland had COVID-19 (95% credible interval: 281,400 to 348,700), equating to 5.98% of the population or around 1 in 17 people.

Because of a shortened publication cycle for this release (to accommodate the 15 to 18 April 2022 bank holiday weekend), analyses relating to the number of new COVID-19 infections (incidence) and sub-regional positivity rates have not been updated this week. Previous estimates of incidence can be found in [our Coronavirus \(COVID-19\) Infection Survey datasets](#). Complete sequenced data for COVID-19 variants have not been updated this week, with most recent data from the week ending 20 March 2022 reported in [Section 6](#).

Within this bulletin, we summarise some of the latest results from the Coronavirus (COVID-19) Infection Survey. For more detailed information on our methods, see [our Coronavirus \(COVID-19\) Infection Survey methodology article](#).

## About this bulletin

In this bulletin, we refer to the number of current COVID-19 infections within the population living in private residential households. We exclude those in hospitals, care homes and/or other communal establishments. In communal establishments, rates of COVID-19 infection are likely to be different. More information about the COVID-19 pandemic from the Office for National Statistics (ONS) and other sources can be found in [our Coronavirus \(COVID-19\) latest insights](#).

The positivity rate is the percentage of people who have tested positive for COVID-19 on a polymerase chain reaction (PCR) test at a point in time. We use current COVID-19 infections to mean testing positive for SARS-CoV-2, with or without having symptoms, on a swab taken from the nose and throat. This is different to the incidence rate, which is a measure of only the new PCR positive cases in a given time period. All analysis was produced with our research partners at the University of Oxford.

Our estimates are based on confirmed positive test results. The remaining swabs are either negative, which are included in our analysis, or are inconclusive, which are not included in our analysis. Some swabs are test failures, which are also not included in our analysis. The impact of excluding inconclusive results on our estimates of positive infections is likely to be very small and unlikely to affect the trend.

### More about coronavirus

- Find the latest on [coronavirus \(COVID-19\) in the UK](#).
- [Explore the latest coronavirus data and analysis](#) from the ONS and other sources.
- View [all coronavirus data](#).
- Find out how we are [working safely in our studies and surveys](#).

[Early management information](#) from the Coronavirus (COVID-19) Infection Survey is made available to government decision-makers to inform their response to COVID-19. Occasionally we may publish figures early if it is considered in the public interest. We will ensure that we pre-announce any ad hoc or early publications as soon as we can. These will include supporting information where possible to aid user understanding. This is consistent with guidance from the Office for Statistics Regulation (OSR).

## How the data in this bulletin can be used

The data can be used for:

- estimating the number of positive cases among the population living in private households, including cases where people do not report having any symptoms
- identifying differences in numbers of positive cases between UK countries and different regions in England
- estimating the number of new cases and change over time in positive cases

The data cannot be used for:

- measuring the number of cases and infections in care homes, hospitals and/or other communal establishments
- providing information about recovery time of those infected

The results in this bulletin are:

- provisional and subject to revision
- based on infections occurring in private households
- subject to uncertainty; [a credible or confidence interval](#) gives an indication of the uncertainty of an estimate from data analysis

## 2 . Percentage of people who had COVID-19 in UK countries

In England and Northern Ireland, the percentage of people testing positive for coronavirus (COVID-19) decreased in the week ending 9 April 2022. In Wales, the percentage of people testing positive for COVID-19 remained high in the week ending 9 April 2022. In Scotland, the percentage of people testing positive for COVID-19 continued to decrease in the week ending 9 April 2022. Our estimates contain Omicron BA.1 and BA.2 variants and all other variants.

All estimates are based on statistical modelling of the trend in rates of positive nose and throat swab results. All estimates are subject to uncertainty given that a sample is only part of the wider population. Additionally, the estimates for the very latest days may change as more test results are received. Therefore, our official estimates for a particular week are based on a [reference day](#) a few days before the end of the week and caution should be taken in over-interpreting small movements in the very latest trends.

Table 1: Official estimates of the percentage of the population testing positive for COVID-19, UK countries  
Estimated percentage of the population testing positive for COVID-19 on nose and throat swabs, UK, 3 April to 9 April 2022

Country	Estimated average % of the population testing positive for COVID-19	95% credible interval		Estimated average number of people testing positive for COVID-19	95% credible interval		Estimated average ratio of the population testing positive for COVID-19	95% credible interval	
		Lower	Upper		Lower	Upper		Lower	Upper
England	6.92	6.72	7.13	3,773,800	3,662,100	3,888,000	1 in 14	1 in 15	1 in 14
Wales	7.63	6.79	8.55	231,900	206,200	259,800	1 in 13	1 in 15	1 in 12
Northern Ireland	5.23	4.30	6.21	95,900	78,800	114,000	1 in 19	1 in 25	1 in 16
Scotland	5.98	5.35	6.62	314,800	281,400	348,700	1 in 17	1 in 19	1 in 15

Source: Office for National Statistics – Coronavirus (COVID-19) Infection Survey

### Notes:

1. The ratios presented are rounded to the nearest 100 if over 1,000, to the nearest 10 if under 1,000, to the nearest 5 if under 100 and to 1 if under 20. This may result in credible intervals that appear to be similar to the estimated average ratio.
2. These ratios do not represent a person's risk of becoming infected, since risk of infection depends on a number of factors including contact with others or vaccination status.

Because of a relatively smaller number of tests in Wales, Northern Ireland and Scotland in the sample, credible intervals are wider, and results should be interpreted with caution. Wide credible intervals mean that differences between the central estimates within and between nations may appear smaller or more exaggerated than they really are.

**Figure 1: The percentage of people testing positive for COVID-19 decreased in England and Northern Ireland, continued to decrease in Scotland, and remained high in Wales in the most recent week**

Estimated percentage of the population testing positive for COVID-19 on nose and throat swabs, UK, 18 April 2021 to 9 April 2022

## Notes:

1. Official reported estimates are plotted at a reference point believed to be most representative of the given week.
2. Official estimates present the best estimate at that point in time. Modelled estimates are used to calculate the official reported estimate. The model smooths the series to understand the trend and is revised each week to incorporate new test results, providing the best indication of trend over time.
3. Official estimates are displayed over a rolling year up to the most recent week. The full time series of our official estimates from 27 April 2020 onwards are available in [our Coronavirus \(COVID-19\) Infection Survey datasets](#).

[Download the data](#)

## About our estimates

Our headline estimates of the percentage of people testing positive in England, Wales, Northern Ireland and Scotland are the latest official estimates. We include different measures to support our estimation and this section outlines the approaches used.

Official estimates should be used to understand the positivity rate for a single point in time. This is based on the modelled estimate for the latest week and is our best and most stable estimate, used in all previous outputs. The modelled estimate is more suited to understanding the recent trend. This is because the model is regularly updated to include new test results and smooths the trend over time. These modelled estimates can be found in [our Coronavirus \(COVID-19\) Infection Survey datasets](#).

The estimates for non-overlapping 14-day periods, which underpin our modelled official estimates, and the unweighted sample counts are also included in the associated datasets. These estimates are produced using a different method of weighting to the model. For more information on our methods and quality surrounding the estimates please see [our Coronavirus \(COVID-19\) Infection Survey methods article](#) and [our Quality and Methodology Information report](#).

All estimates presented in this bulletin are provisional results. As swabs are not necessarily analysed in date order by the laboratory, we have not yet received test results for all swabs taken on the dates included in this analysis. Estimates may therefore be revised as more test results are included.

## 3 . Sub-national analysis of the percentage of people who had COVID-19

In the week ending 9 April 2022, the percentage of people testing positive for coronavirus (COVID-19) decreased in the North West, East of England, London, South East and South West. The percentage of people testing positive continued to increase in the North East in the week ending 9 April 2022. The percentage of people testing positive increased in Yorkshire and The Humber, East Midlands and West Midlands over the most two recent weeks. However, the trend is uncertain in the most recent week (up to 9 April 2022).

In the data used to produce these estimates, the number of people sampled in each region who tested positive for COVID-19 was low relative to England overall. This means there is a higher degree of uncertainty in the regional estimates, as indicated by larger credible intervals.

### Figure 2: The trend in the percentage of people testing positive for COVID-19 varied across regions in England in the week ending 9 April 2022

Modelled daily percentage of the population testing positive for COVID-19 on nose and throat swabs by region, England, 27 February to 9 April 2022

## Notes:

1. Credible intervals widen slightly at the end as there is a delay between the swab being taken and reporting of results. We report latest figures based on the reference day for that week because of this greater uncertainty in the most recent days.
2. The percentage of people testing positive by region was calculated using a similar modelling approach to the national daily estimates in [Section 2: Percentage of people who had COVID-19 in England, Wales, Northern Ireland and Scotland](#).
3. The analysis is conducted over a six-week period, which means some positive cases move in and out of the sample. This causes variability between estimates over time, which is expected given the lower number of positive tests in each region, compared with England as a whole.
4. We describe trends by comparing the probability that the estimate for the reference day is higher or lower than the estimate for 7 and 14 days prior.

[Download the data](#)

Estimates for non-overlapping 14-day periods (which underpin our modelled estimates) by region are available in [our Coronavirus \(COVID-19\) Infection Survey datasets](#).

## Sub-regional analysis of the UK

Because of a shortened publication cycle for this release (to accommodate the 15 to 18 April 2022 bank holiday weekend), sub-regional analyses have not been updated this week. Our latest estimates for week ending 2nd April can be found in [our Coronavirus \(COVID-19\) Infection Survey datasets](#).

# 4 . Age analysis of the percentage of people who had COVID-19

## Age group analysis for England

Our [age groups](#) separate children and young people by school age.

In the week ending 9 April 2022, the percentage of people testing positive decreased in all age groups except those in School Year 7 to School Year 11 and those aged 70 years and over for which the trend was uncertain. The percentage of people testing positive increased for those aged 70 years and over in the most recent two weeks, but the trend is uncertain in the most recent week (up to 9 April 2022). Caution should be taken in over-interpreting any small movements in the latest trend.

**Figure 3: The percentage of people testing positive for COVID-19 in England decreased in all age groups except those in School Year 7 to School Year 11 and those aged 70 years and over for which the trend was uncertain in the most recent week up to 9 April 2022**

Modelled daily percentage of the population testing positive for COVID-19 on nose and throat swabs by age group, England, 27 February to 9 April 2022

Notes:

1. Credible intervals widen slightly at the end as there can be a delay between the swab being taken and reporting of results. We report latest figures based on the reference day for that week because of this greater uncertainty in the most recent days.

[Download the data](#)

Estimates are based on smaller sample sizes within each age group relative to England overall. There is a higher degree of uncertainty as indicated by larger credible intervals. These can be found in the [Coronavirus \(COVID-19\) Infection Survey datasets](#).

Estimates for non-overlapping 14-day periods (which underpin our modelled estimates) by age are available in [our Coronavirus \(COVID-19\) Infection Survey datasets](#).

We are unable to produce the same grouped analysis as presented in Figure 3 for the devolved administrations because of smaller sample sizes within each age group. However, estimates on positivity by single year of age for Wales, Northern Ireland and Scotland using a different model are in the following section and [our accompanying datasets](#).

## Age analysis by single year of age over time by country

In this section, we present modelled daily estimates of the percentage testing positive for COVID-19 by single year of age over time from 27 February to 9 April 2022 for England, Wales, Northern Ireland and Scotland. They are produced using a different method to the grouped age analysis for England presented previously and are therefore not comparable.

Estimates are based on smaller sample sizes within each age group in comparison with the total sample size for each country. There is a higher degree of uncertainty as indicated by larger confidence intervals. These can be found in [our Coronavirus \(COVID-19\) Infection Survey datasets](#).

The data presented in Figure 4 for England suggest that the percentage of people testing positive decreased in all ages except for those aged 70 years and over.

For Wales, rates have generally increased for those aged around 40 years and over while the trend is uncertain among younger adults and children. For Northern Ireland, rates have generally decreased in recent weeks for younger adults and children, while the trend is uncertain for older adults. For Scotland, rates have generally decreased among all ages in recent weeks.

Further analysis on age for [Wales](#), [Northern Ireland](#) and [Scotland](#) is published by their respective statistical agencies. Analysis for Wales is published in [English](#) and [Welsh](#).

### Figure 4: The percentage of people testing positive for COVID-19 by single year of age over time for England, Wales, Northern Ireland and Scotland

Modelled daily percentage of the population testing positive for COVID-19 on nose and throat swabs by single year of age, UK, 27 February to 9 April 2022

#### Notes:

1. These estimates use a different method to modelled daily estimates of the percentage testing positive by age group for England in the previous section and are therefore not comparable.

[Download the data](#)

## 5 . Number of new COVID-19 infections in UK countries

Because of a shortened publication cycle for this release (to accommodate the 15 to 18 April 2022 bank holiday weekend), analyses relating to the number of new COVID-19 infections (incidence) have not been updated this week. Previous estimates of incidence can be found in [our Coronavirus \(COVID-19\) Infection Survey datasets](#).

## 6 . Analysis of viral load and variants of COVID-19

We publish [cycle-threshold \(Ct\) values](#) and a breakdown of infections by variant, based on gene positivity patterns from our swab tests. The "Cycle threshold", known as a Ct value, is a proxy for the quantity of virus (also known as viral load), where a lower Ct value indicates higher viral load. The latest Ct values of coronavirus (COVID-19) positive tests are provided in [our Coronavirus \(COVID-19\) Infection Survey: technical dataset](#).

The Omicron variant BA.1 has changes in one of the three genes that the coronavirus survey swab test detects, which means the S-gene is no longer detected. When there is a high viral load (for example, when a person is most infectious), not detecting the S-gene in combination with detecting the other two genes (ORF1ab and N-genes) is a reliable indicator of this Omicron BA.1 variant. However, as the viral load decreases (for example, if someone is near the end of their recovery from the infection), not detecting the S-gene is a less reliable indicator of this Omicron variant.

The Omicron variant BA.2 does not have changes in the S-gene, and therefore all three genes, or the S-gene and either ORF1ab or N, will usually be detected in infections with this variant.

Our latest complete sequenced data were last updated the week ending 20 March 2022 and showed that the Omicron BA.2 variant remained the most common variant in England, Wales, Northern Ireland and Scotland. In that week, 89.8% of all sequenced COVID-19 infections from the survey were compatible with the Omicron BA.2 variant, and 10.2% were compatible with the Omicron BA.1 variant or its sub-variants.

Our main variant analysis is for a reference day, and therefore is not directly comparable with the sequencing data. The following analysis looks at the percentage of the population with a positive test compatible with the Omicron BA.1 or BA.2 variant in England, Wales, Northern Ireland and Scotland, and the regions of England.

During periods of change in COVID-19 variants we will include a breakdown of estimated infections by variant to illustrate how the estimated percentage of people infected in the population is changing by variant (Figure 5). When nearly all infections are compatible with a dominant variant, we will no longer include this breakdown.

Data should be treated with caution. There could be small numbers of positive cases compatible with a particular variant in Wales, Scotland and Northern Ireland, leading to considerable uncertainty surrounding these estimates. Not all cases that are positive on the ORF1ab and N-genes will be the Omicron BA.1 variant, and not all cases that have a detectable S-gene will be the Omicron BA.2 variant.

In the week up to 9 April 2022, the percentage of people with infections compatible with the Omicron BA.2 variant decreased in England, Northern Ireland and Scotland. The percentage of people with infections compatible with the Omicron BA.2 variant remained high in Wales in the most recent week (up to 9 April 2022).

In the same week, the percentage of people with infections compatible with the Omicron BA.1 variant continued to decrease in England and Scotland, while the trend was uncertain for Wales and Northern Ireland.

### **Figure 5: The percentage of people with infections compatible with the Omicron BA.2 variant decreased in England, Northern Ireland and Scotland, and remained high in Wales in the most recent week**

**Modelled percentage of the population testing positive for infections compatible with the Omicron BA.1 variant and Omicron BA.2 variant, based on nose and throat swabs, daily, UK, 27 February to 9 April 2022**

**Notes:**

1. Omicron BA.1 variant-compatible positives are defined as those that are positive on the ORF1ab-gene and N-gene, but not the S-gene. This group includes Omicron BA.1.1.
2. Omicron BA.2 variant-compatible positives are defined as those that are positive on the ORF1ab-gene, N-gene and S-gene, on the ORF1ab-gene and S-gene, or on the N-gene and S-gene.
3. Data should be treated with caution. Not all infections positive on the S-gene will be the Omicron BA.2 variant, and some infections with pattern ORF1ab plus N will also be the Omicron BA.2 variant where the S-gene was not detected for other reasons, such as low viral load.

### [Download the data](#)

In the week ending 9 April 2022, the percentage of people testing positive with infections compatible with the Omicron BA.2 variant decreased in the North West, East of England, London, South East and South West. The percentage of people testing positive with infections compatible with the Omicron BA.2 variant increased in the most recent week in the North East. The trend is uncertain in Yorkshire and The Humber, East Midlands and West Midlands in the most recent week (up to 9 April 2022).

In the same week, the percentage of people with infections compatible with the Omicron BA.1 variant decreased in the East Midlands, East of England, London, South East and South West, while the trend was uncertain in the North East, North West, Yorkshire and The Humber and West Midlands. These modelled estimates can be found in [our Coronavirus \(COVID-19\) Infection Survey datasets](#).

The [World Health Organization \(WHO\) have defined names for variants of concern](#). These are variants that the UK government has under surveillance. You can find out more in the latest [SARS-CoV-2 variants of concern and variants under investigation in England briefing document on GOV.UK](#).

Currently circulating variants of concern in the UK are:

- Delta: B.1.617.2 and its genetic descendants
- Omicron: B.1.1.529 (which includes sublineages BA.1, BA.1.1, BA.2 and BA.3)

[Our Coronavirus \(COVID-19\) Infection Survey: technical dataset](#) includes analysis of the genetic lineages of coronavirus seen in the samples we sequence. More information on how we measure variants from positive tests on the survey can be found in [our Understanding COVID-19 variants blog](#). [Our Coronavirus \(COVID-19\) Infection Survey methods article](#) gives more detail about how we sequence the virus' genetic material.

The sequencing is produced by Northumbria University and analysis is produced by research partners at the University of Oxford. Of particular note are Dr Katrina Lythgoe, Dr Tanya Golubchik and Dr Helen Fryer. Genome sequencing is funded by the COVID-19 Genomics UK (COG-UK) consortium. COG-UK is supported by funding from the Medical Research Council (MRC) part of UK Research and Innovation (UKRI), the National Institute of Health Research (NIHR), and Genome Research Limited operating as the Wellcome Sanger Institute.

## 7 . Test sensitivity and specificity

The estimates provided in Sections 2 to 6 are for the percentage of the private-residential population testing positive for coronavirus (COVID-19), otherwise known as the positivity rate. We do not report the prevalence rate. To calculate the prevalence rate, we would need an accurate understanding of the swab test's sensitivity (true-positive rate) and specificity (true-negative rate).

While we do not know the true sensitivity and specificity of the test, our data and related studies provide an indication of what these are likely to be. In particular, the data suggest that the false-positive rate is very low - under 0.005%. We do not know the sensitivity of the swab test. However, other studies suggest that sensitivity (the rate of true-positive test results) may be somewhere between 85% and 98%.

You can find more information on sensitivity and specificity in [our Coronavirus \(COVID-19\) Infection Survey methods article](#) and [our blog that explains why we trust the data from the Coronavirus \(COVID-19\) Infection Survey](#). You can find [more information on the data suggesting that our test's false-positive rate is very low](#) in a paper written by academic partners at the University of Oxford.

## 8 . Coronavirus (COVID-19) Infection Survey data

### [Coronavirus \(COVID-19\) Infection Survey: England](#)

Dataset | Released 14 April 2022

Findings from the Coronavirus (COVID-19) Infection Survey for England.

### [Coronavirus \(COVID-19\) Infection Survey: Northern Ireland](#)

Dataset | Released 14 April 2022

Findings from the Coronavirus (COVID-19) Infection Survey for Northern Ireland.

### [Coronavirus \(COVID-19\) Infection Survey: Scotland](#)

Dataset | Released 14 April 2022

Findings from the Coronavirus (COVID-19) Infection Survey for Scotland.

### [Coronavirus \(COVID-19\) Infection Survey: Wales](#)

Dataset | Released 14 April 2022

Findings from the Coronavirus (COVID-19) Infection Survey for Wales.

### [Coronavirus \(COVID-19\) Infection Survey: technical data](#)

Dataset | Released 14 April 2022

Technical and methodological data from the Coronavirus (COVID-19) Infection Survey, England, Wales, Northern Ireland and Scotland.

## 9 . Collaboration



The Coronavirus (COVID-19) Infection Survey analysis was produced by the Office for National Statistics (ONS) in collaboration with our research partners at the University of Oxford, the University of Manchester, UK Health Security Agency (UK HSA) and Wellcome Trust. Of particular note are:

- Sarah Walker - University of Oxford, Nuffield Department for Medicine: Professor of Medical Statistics and Epidemiology and Study Chief Investigator
- Koen Pouwels - University of Oxford, Health Economics Research Centre, Nuffield Department of Population Health: Senior Researcher in Biostatistics and Health Economics
- Thomas House - University of Manchester, Department of Mathematics: Reader in Mathematical Statistics
- Anna Seale - University of Warwick, Warwick Medical School: Professor of Public Health; UK Health Security Agency, Data, Analytics and Surveillance: Scientific Advisor

## 10 . Glossary

### Age groups for children and young people

- "aged 2 years to school Year 6" includes children in primary school and below
- "school Year 7 to school Year 11" includes children in secondary school
- "school Year 12 to those aged 24 years" includes young adults who may be in further or higher education

Those aged from 11 to 12 years and those aged 16 to 17 years have been split between different age categories depending on whether their birthday is before or after 1 September.

### Confidence interval

A confidence interval gives an indication of the degree of uncertainty of an estimate, showing the precision of a sample estimate. The 95% confidence intervals are calculated so that if we repeated the study many times, 95% of the time the true unknown value would lie between the lower and upper confidence limits. A wider interval indicates more uncertainty in the estimate. Overlapping confidence intervals indicate that there may not be a true difference between two estimates. For more information, see [our Statistical uncertainty methodology page](#).

### Credible interval

A credible interval gives an indication of the uncertainty of an estimate from data analysis. The 95% credible intervals are calculated so that there is a 95% probability of the true value lying in the interval.

### Cycle threshold (Ct) values

The strength of a positive coronavirus (COVID-19) test is determined by how quickly the virus is detected, measured by a cycle threshold (Ct) value. The lower the Ct value, the higher the viral load and stronger the positive test. Positive results with a high Ct value can be seen in the early stages of infection when virus levels are rising, or late in the infection, when the risk of transmission is low.

### False-positives and false-negatives

A false-positive result occurs when the tests suggest a person has COVID-19 when in fact they do not. By contrast, a false-negative result occurs when the tests suggest a person does not have COVID-19 when in fact they do. For more information on false-positives and false-negatives, see our [our Coronavirus Infection Survey methods article](#) and [our Accuracy and confidence: why we trust the data from the COVID-19 infection survey blog](#).

### Incidence rate

The incidence rate is a measure of the estimated number of new polymerase chain reaction (PCR)-positive cases per day per 10,000 people at a given point in time. It is different to positivity, which is an estimate of all current PCR positive cases at a point in time, regardless of whether the infection is new or existing.

# 11 . Measuring the data

## Reference dates

We aim to provide the estimates of positivity rate (the percentage of those who test positive) and incidence that are most timely and most representative of each week. We decide the most recent week we can report on based on the availability of test results for visits that have already happened, accounting for the fact that swabs have to be couriered to the labs, tested and results returned. On most occasions, the reference dates align perfectly, but sometimes this is not feasible. This week, the reference week for positivity is 3 to 9 April 2022 for England, Wales, Northern Ireland and Scotland.

Within the most recent week, we provide an official estimate for positivity rate based on a reference point from the modelled trends. For positivity rates, we can include all swab test results, even from the most recent visits. Therefore, although we are still expecting further swab test results from the labs, there were sufficient data for the official estimate for infection to be based on a reference point after the start of the reference week. To improve stability in our modelling while maintaining relative timeliness of our estimates, we are reporting our official estimates based on the midpoint of the reference week. This week, the reference day for positivity rates is Wednesday 6 April 2022 for England, Wales, Northern Ireland and Scotland.

The reference date used for our official estimates of incidence of polymerase chain reaction (PCR)-positive cases is 14 days prior to the positivity reference day. This is necessary as estimates later than this date are more likely to change as we receive additional data. The most recent analyses of incidence are not included in this bulletin; the reference week for incidence in the previous release of this bulletin (8 April 2022) was 13 to 19 March 2022 and the reference day was Wednesday 16 March 2022 for all UK countries.

## Response rates

Enrolment for this wave of recruitment for the Coronavirus (COVID-19) Infection Survey ceased on 31 January 2022. Response rates for England, Wales, Northern Ireland and Scotland can be regarded as final response rates to the survey. Response rates for each nation are found in [our Coronavirus \(COVID-19\) Infection Survey: technical dataset](#). We provide response rates separately for the different sampling phases of the study. Additional information on response rates can be found in [our Coronavirus \(COVID-19\) Infection Survey methods article](#).

## Survey fieldwork

Survey fieldwork for the pilot study began in England on 26 April 2020. In Wales, fieldwork began on 29 June 2020, in Northern Ireland fieldwork began on 26 July 2020 and in Scotland fieldwork began on 21 September 2020.

## Sub-regional geographies

We have presented modelled estimates for the most recent week of data at the sub-regional level. To balance granularity with statistical power, we have grouped together local authorities into Coronavirus (COVID-19) Infection Survey sub-regions. The geographies are a rules-based composition of local authorities. Local authorities with a population over 200,000 have been retained where possible.

The boundaries for these Coronavirus (COVID-19) Infection Survey sub-regions can be found on the [Open Geography Portal](#).

## Other Coronavirus Infection Survey (CIS) analysis and studies

This study is one of a number of studies that look to provide information around the coronavirus pandemic within the UK. For information on other studies see [Section 5: Quality characteristics of the Coronavirus \(COVID-19\) Infection Survey](#) (coherence and comparability), revised 16 July 2021.

## 12 . Strengths and limitations

These statistics have been produced quickly in response to developing world events. The [Office for Statistics Regulation \(OSR\)](#), on behalf of the UK Statistics Authority, has reviewed them on [14 May 2020](#) and [17 March 2021](#) against several important aspects of the [Code of Practice for Statistics](#) and regards them as consistent with the Code's pillars of [trustworthiness](#), [quality](#) and [value](#).

The estimates presented in this bulletin contain [uncertainty](#). There are many sources of uncertainty, including uncertainty in the test, in the estimates and in the quality of data collected in the questionnaire. Information on the main sources of uncertainty are presented in [our Coronavirus \(COVID-19\) Infection Survey Quality and Methodology Information report](#), [our Coronavirus \(COVID-19\) Infection Survey: methods and further information article](#), and [our blog that explains why we trust the data from the Coronavirus \(COVID-19\) Infection Survey](#).

## 13 . Related links

### [Coronavirus \(COVID-19\) Infection Survey: characteristics of people testing positive for COVID-19 in countries of the UK](#)

Bulletin | Updated fortnightly

The characteristics of people testing positive for coronavirus (COVID-19) from the COVID-19 Infection Survey. This survey is being delivered in partnership with the University of Oxford, the University of Manchester, UK Health Security Agency and Wellcome Trust.

### [Coronavirus \(COVID-19\) Infection Survey: antibody and vaccination data for the UK](#)

Bulletin | Updated fortnightly

Antibody and vaccination data by UK country and English regions from the Coronavirus (COVID-19) Infection Survey. This survey is being delivered in partnership with the University of Oxford, University of Manchester, UK Health Security Agency and Wellcome Trust.

### [Coronavirus \(COVID-19\) latest insights](#)

Interactive tool | Updated as and when data become available

The latest data and trends about the coronavirus (COVID-19) pandemic from the Office for National Statistics (ONS) and other official sources.

### [Deaths registered weekly in England and Wales, provisional](#)

Bulletin | Updated weekly

Provisional counts of the number of deaths registered in England and Wales, including deaths involving COVID-19, by age, sex and region, in the latest weeks for which data are available.

### [Coronavirus \(COVID-19\) Infection Survey technical article: Analysis of characteristics associated with vaccination uptake](#)

Technical article | Released 15 November 2021

Analysis of populations in the UK by likelihood of being vaccinated against COVID-19 using the Coronavirus (COVID-19) Infection Survey. This survey is being delivered in partnership with University of Oxford, University of Manchester, UK Health Security Agency and Wellcome Trust.

### [Coronavirus \(COVID-19\) Infection Survey: methods and further information](#)

Methodology article | Updated 7 February 2022

Information on the methods used to collect the data, process it, and calculate the statistics produced from the Coronavirus (COVID-19) Infection Survey.

### [The relationship between COVID-19 infections and antibodies: What do the data show?](#)

Blog article | Released 6 April 2022

Information on how COVID-19 infections have recently reached their highest levels across many parts of the UK while antibody levels are high.

### [Coronavirus \(COVID-19\) Infection Survey QMI](#)

Methodology article | Updated 16 July 2021

Quality and Methodology Information for the Coronavirus (COVID-19) Infection Survey (CIS), detailing the strengths and limitations of the data, methods used, and data uses and users.